Affymetrix. (2001). Affymetrix ® Microarray Suite User's Guide (Santa Clara: Affymetrix).

Ager, F.J., Ynsa, M.D., Dominguez-Solis, J.R., Gotor, C., Respaldiza, M.A., Romero, L.C. (2002). Cadmium localization and quantification in the plant *Arabidopsis thaliana* using micro-PIXE. Nucl. Instrum. Methods Phys. Res. Sect. B-Beam Interact. Mater. Atoms **189,** 494-498.

Agrawal, G.K., Agrawal, S.K., Shibato, J., Iwahashi, H., Rakwal, R. (2003). Novel rice MAP kinases *OsMSRMK3* and *OsWJUMK1* involved in encountering diverse environmental stresses and developmental regulation. Biochem. Biophys. Res. Commun. **300,** 775-783.

Agrawal, G.K., Rakwal, R., Iwahashi, H. (2002). Isolation of novel rice (*Oryza sativa* L.) multiple stress responsive MAP kinase gene, *OsMSRMK2*, whose mRNA accumulates rapidly in response to environmental cues. Biochem. Biophys. Res. Commun. **294,** 1009-1016.

Allen, G.J., Chu, S.P., Harrington, C.L., Schumacher, K., Hoffmann, T., Tang, Y.Y., Grill, E., Schroeder, J.I. (2001). A defined range of guard cell calcium oscillation parameters encodes stomatal movements. Nature **411,** 1053-1057.

Allen, G.J., Kwak, J.M., Chu, S.P., Llopis, J., Tsien, R.Y., Harper, J.F., Schroeder, J.I. (1999). Cameleon calcium indicator reports cytoplasmic calcium dynamics in *Arabidopsis* guard cells. Plant J. **19,** 735-747

Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J. (1990). Basic local alignment search tool. J. Mol. Biol. **215,** 403-410.

Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W., Lipman, D.J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucl. Acids Res. **25,** 3389-3402.

Anderson, C.W.N., Brooks, R.R., Stewart, R.B., Simcock, R. (1998). Harvesting a crop of gold in plants. Nature **395,** 553-554.

Andrews, N.C. (2002). Metal transporters and disease. Curr. Opin. Chem. Biol. **6,** 181-186.

Anton, A., Grosse, C., Reissmann, J., Pribyl, T., Nies, D.H. (1999). CzcD is a heavy metal ion transporter involved in regulation of heavy metal resistance in *Ralstonia* sp strain CH34. J. Bacteriol. **181,** 6876-6881.

Arazi, T., Sunkar, R., Kaplan, B., Fromm, H. (1999). A tobacco plasma membrane calmodulin-binding transporter confers Ni^{2+} tolerance and Pb^{2+} hypersensitivity in transgenic plants. Plant J. **20,** 171-182.

Arner, E.S., Holmgren, A. (2000). Physiological functions of thioredoxin and thioredoxin reductase. Eur. J. Biochem. **267,** 6102-6109.

Asai, T., Tena, G., Plotnikova, J., Willmann, M.R., Chiu, W.L., Gomez-Gomez, L., Boller, T., Ausubel, F.M., Sheen, J. (2002). MAP kinase signalling cascade in *Arabidopsis* innate immunity. Nature **415,** 977-983.

Assuncao, A.G.L., Martins, P.D., De Folter, S., Vooijs, R., Schat, H., Aarts, M.G.M. (2001). Elevated expression of metal transporter genes in three accessions of the metal hyperaccumulator *Thlaspi caerulescens*. Plant Cell Environ. **24,** 217-226.

Ausubel, F.M., Brent, R., Kingston, R.E., Moore, D.D., Seidman, J.G., Smith, J.A., Struhl, K. (1989). Current Protocols in Molecular Biology. (New York: John Wiley and Sons, Inc.).

Avsian-Kretchmer, O., Eshdat, Y., Gueta-Dahan, Y., Ben-Hayyim, G. (1999). Regulation of stress-induced phosholipid hydroperoxide glutathione peroxidase expression in citrus. Planta **209,** 469-477.

Bachem, C.W., van der Hoeven, R.S., de Bruijn, S.M., Vreugdenhil, D., Zabeau, M., Visser, R.G. (1996). Visualization of differential gene expression using a novel method of RNA fingerprinting based on AFLP: analysis of gene expression during potato tuber development. Plant J. **9,** 745-753.

Bailey, T.L., Elkan, C. (1994). Fitting a mixture model by expectation maximization to discover motifs in biopolymers. In Proceedings of Second International Conference on Intelligent Systems for Molecular Biology (Melno Park, California: AAAI Press), 28-36.

Bailey, T.L., Gribskov, M. (1998). Combining evidence using p-values: application to sequence homology searches. Bioinformatics **14,** 48-54.

Baker, A.J.M., Brooks, R.R. (1989). Terrestrial Higher Plants which Hyperaccumulate Metallic Elements - A review of their Distribution, Ecology and Phytochemistry. Biorecovery **1,** 81-126.

Baker, A.J.M., McGrath, S.P., Reeves, R.D., Smith, J.A.C. (2000). Metal hyperaccumulator plants: a review of the ecology and physiology of a biochemical resource for phytoremediation of metal-polluted soils. In Phytoremediation of contaminated soil and water. N. Terry, G. Banuelos, and J. Vangronsveld, (Boca-Raton, FL, USA: Lewis Publishers), 85-107.

Balandin, T., Castresana, C. (2002). AtCOX17, an *Arabidopsis* homolog of the yeast copper chaperone COX17. Plant Physiol. **129,** 1852-1857.

Baldwin, D., Crane, V., Rice, D. (1999). A comparison of gel-based, nylon filter and microarray techniques to detect differential RNA expression in plants. Curr. Opin. Plant Biol. **2,** 96-103.

Barcelo, J., Poschenrieder, C. (1990). Plant Water Relations as Affected by Heavy-Metal Stress - a Review. J. Plant Nutr. **13,** 1-37.

Baudouin, E., Meskiene, I., Hirt, H. (1999). Unsaturated fatty acids inhibit MP2C, a protein phosphatase 2C involved in the wound-induced MAP kinase pathway regulation. Plant J. **20,** 343-348.

Becraft, P.W. (2002). Receptor Kinase Signaling in Plant Development. Annu. Rev. Cell Dev. Biol. **18,** 163-192.

Ben-Hayyim, G., Gueta-Dahan, Y., Avsian-Kretchmer, O., Weichert, H., Feussner, I. (2001). Preferential induction of a 9-lipoxygenase by salt in salt-tolerant cells of *Citrus sinensis* L. Osbeck. Planta **212,** 367-375.

Bert, V., Bonnin, I., Saumitou-Laprade, P., de Laguerie, P., Petit, D. (2002). Do *Arabidopsis halleri* from nonmetallicolous populations accumulate zinc and cadmium more effectively than those from metallicolous populations? New Phytol. **155,** 47-57.

Bert, V., Macnair, M.R., De Laguerie, P., Saumitou-Laprade, P., Petit, D. (2000). Zinc tolerance and accumulation in metallicolous and nonmetallicolous populations of *Arabidopsis halleri (*Brassicaceae). New Phytol. **146,** 225-233.

Bert, V., Meerts, P., Saumitou-Laprade, P., Salis, P., Gruber, W., Verbruggen, N. (2003). Genetic basis of Cd tolerance and hyperaccumulation in *Arabidopsis halleri*. Plant Soil **249,** 9-18.

Birnboim, H.C., Doly, J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. Nucl. Acids Res. **7,** 1513-1523.

Bizily, S.P., Rugh, C.L., Meagher, R.B. (2000). Phytodetoxification of hazardous organomercurials by genetically engineered plants. Nat. Biotechnol. **18,** 213-217.

Bizily, S.P., Rugh, C.L., Summers, A.O., Meagher, R.B. (1999). Phytoremediation of methylmercury pollution: merB expression in *Arabidopsis thaliana* confers resistance to organomercurials. Proc. Natl. Acad. Sci. USA **96,** 6808-6813.

Bloss, T., Clemens, S., Nies, D.H. (2002). Characterization of the ZAT1p zinc transporter from *Arabidopsis thaliana* in microbial model organisms and reconstituted proteoliposomes. Planta **214,** 783-791.

Bogre, L., Calderini, O., Binarova, P., Mattauch, M., Till, S., Kiegerl, S., Jonak, C., Pollaschek, C., Barker, P., Huskisson, N.S., Hirt, H., Heberle-Bors, E. (1999). A MAP kinase is activated late in plant mitosis and becomes localized to the plane of cell division. Plant Cell **11,** 101-113.

Bogre, L., Ligterink, W., Heberle-Bors, E., Hirt, H. (1996). Mechanosensors in plants. Nature **383,** 489-490.

Bogre, L., Ligterink, W., Meskiene, I., Barker, P.J., Heberle-Bors, E., Huskisson, N.S., Hirt, H. (1997). Wounding Induces the Rapid and Transient Activation of a Specific MAP Kinase Pathway. Plant Cell **9,** 75-83.

Bovet, L., Eggmann, T., Meylan-Bettex, M., Polier, J.E., Kammer, P., Marin, E., Feller, U., Martinoia, E. (2003). Transcript levels of AtMRPs after cadmium treatment: induction of *AtMRP3*. Plant Cell Environ. **26,** 371-381.

Breyne, P., Dreesen, R., Cannoot, B., Rombaut, D., Vandepoele, K., Rombauts, S., Vanderhaeghen, R., Inze, D., Zabeau, M. (2003). Quantitative cDNA-AFLP analysis for genome-wide expression studies. Mol. Genet. Genomics **269,** 173-179.

Brooks, R.R., Chambers, M.F., Nicks, L.J., Robinson, B.H. (1998). Phytomining. Trends Plant Sci. **3,** 359-362.

Buchanan, B.B., Gruissem, W., Jones, R.L. (2000). Biochemistry & Molecular Biology of Plants. (Rockville, Maryland: American Society of Plant Physiologists).

Butt, A., Mousley, C., Morris, K., Beynon, J., Can, C., Holub, E., Greenberg, J.T., Buchanan-Wollaston, V. (1998). Differential expression of a senescence-enhanced metallothionein gene in *Arabidopsis* in response to isolates of *Peronospora parasitica* and *Pseudomonas syringae*. Plant J. **16,** 209-221.

Cardinale, F., Jonak, C., Ligterink, W., Niehaus, K., Boller, T., Hirt, H. (2000). Differential activation of four specific MAPK pathways by distinct elicitors. J. Biol. Chem. **275,** 36734-36740.

Carninci, P., Hayashizaki, Y. (1999). High-Efficiency Full-Length cDNA Cloning. Methods Enzymol. **303,** 19-44.

Carri, M.T., Galiazzo, F., Ciriolo, M.R., Rotilio, G. (1991). Evidence for co-regulation of Cu,Zn superoxide dismutase and metallothionein gene expression in yeast through transcriptional control by copper via the ACE 1 factor. FEBS Lett. **278,** 263-266.

Chaney, R.L. (1988). Plants can utilize iron from Fe-N,N'-di-(2-hydroxybenzoly)-ethylenediamine-N-N'-diacetic acid, a ferric chelate with greater formation constant than Fe-EDDHA. J. Plant Nutr. **11,** 1033-1050.

Chaney, R.L., Malik, M., Li, Y.M., Brown, S.L., Brewer, E.P., Angle, J.S., Baker, A.J. (1997). Phytoremediation of soil metals. Curr. Opin. Biotechnol. **8,** 279-284.

Chardonnens, A.N., Koevoets, P.L., van Zanten, A., Schat, H., Verkleij, J.A. (1999). Properties of enhanced tonoplast zinc transport in naturally selected zinc-tolerant *Silene vulgaris*. Plant Physiol. **120,** 779-786.

Chen, W., Provart, N.J., Glazebrook, J., Katagiri, F., Chang, H.S., Eulgem, T., Mauch, F., Luan, S., Zou, G., Whitham, S.A., Budworth, P.R., Tao, Y., Xie, Z., Chen, X., Lam, S., Kreps, J.A., Harper, J.F., Si-Ammour, A., Mauch-Mani, B., Heinlein, M., Kobayashi, K., Hohn, T., Dangl, J.L., Wang, X., Zhu, T. (2002). Expression profile matrix of *Arabidopsis* transcription factor genes suggests their putative functions in response to environmental stresses. Plant Cell **14,** 559-574.

Chen, W.Q., Chao, G., Singh, K.B. (1996). The promoter of a H_2O_2-inducible, *Arabidopsis* glutathione S-transferase gene contains closely linked OBF- and OBP1-binding sites. Plant J. **10,** 955-966.

Cheng, S.H., Willmann, M.R., Chen, H.C., Sheen, J. (2002). Calcium signaling through protein kinases. The *Arabidopsis* calcium-dependent protein kinase gene family. Plant Physiol. **129,** 469-485.

Cheong, Y.H., Chang, H.S., Gupta, R., Wang, X., Zhu, T., Luan, S. (2002). Transcriptional profiling reveals novel interactions between wounding, pathogen, abiotic stress, and hormonal responses in *Arabidopsis*. Plant Physiol. **129**, 661-77.

Cheung, W.Y. (1988). Calmodulin and its activation by cadmium ion. Ann. N. Y. Acad. Sci. **522,** 74-87.

Chomczynski, P., Sacchi, N. (1987). Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. Anal. Biochem. **162,** 156-159.

Chuang, C.F., Running, M.P., Williams, R.W., Meyerowitz, E.M. (1999). The PERIANTHIA gene encodes a bZIP protein involved in the determination of floral organ number in *Arabidopsis thaliana*. Genes Dev. **13,** 334-344.

Clark, G.B., Sessions, A., Eastburn, D.J., Roux, S.J. (2001). Differential expression of members of the annexin multigene family in *Arabidopsis*. Plant Physiol. **126,** 1072-1084.

Clemens, S. (2001). Molecular mechanisms of plant metal tolerance and homeostasis. Planta **212,** 475-486.

Clemens, S., Antosiewicz, D.M., Ward, J.M., Schachtman, D.P., Schroeder, J.I. (1998). The plant cDNA LCT1 mediates the uptake of calcium and cadmium in yeast. Proc. Natl. Acad. Sci. USA **95,** 12043-12048.

Clemens, S., Bloss, T., Vess, C., Neumann, D., Nies, D.H., zur Nieden, U. (2002b). A transporter in the endoplasmic reticulum of *Schizosaccharomyces pombe* cells mediates zinc storage and differentially affects transition metal tolerance. J. Biol. Chem. **277,** 18215-18221.

Clemens, S., Kim, E.J., Neumann, D., Schroeder, J.I. (1999). Tolerance to toxic metals by a gene family of phytochelatin synthases from plants and yeast. Embo J. **18,** 3325-3333.

Clemens, S., Palmgren, M.G., Kramer, U. (2002a). A long way ahead: understanding and engineering plant metal accumulation. Trends Plant Sci. **7,** 309-315.

Clemens, S., Schroeder, J.I., Degenkolb, T. (2001). *Caenorhabditis elegans* expresses a functional phytochelatin synthase. Eur. J. Biochem. **268,** 3640-3643.

Clough, S.J., Bent, A.F. (1998). Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. Plant J. **16,** 735-743.

Cobbett, C., Goldsbrough, P. (2002). Phytochelatins and metallothioneins: roles in heavy metal detoxification and homeostasis. Annu. Rev. Plant Biol. **53,** 159-182.

Cohen, P. (1989). The Structure and Regulation of Protein Phosphatases. Annu. Rev. Biochem. **58,** 453-508.

Coleman, J.E. (1992). Zinc proteins: enzymes, storage proteins, transcription factors, and replication proteins. Annu. Rev. Biochem. **61,** 897-946.

Conklin, D.S., McMaster, J.A., Culbertson, M.R., Kung, C. (1992). *Cot1*, a Gene Involved in Cobalt Accumulation in *Saccharomyces cerevisiae*. Mol. Cell Biol. **12,** 3678-3688.

Connolly, E.L., Fett, J.P., Guerinot, M.L. (2002). Expression of the *IRT1* metal transporter is controlled by metals at the levels of transcript and protein accumulation. Plant Cell **14,** 1347-1357.

Culotta, V.C., Klomp, L.W.J., Strain, J., Casareno, R.L.B., Krems, B., Gitlin, J.D. (1997). The copper chaperone for superoxide dismutase. J. Biol. Chem. **272,** 23469-23472.

Curie, C., Alonso, J.M., Le Jean, M., Ecker, J.R., Briat, J.F. (2000). Involvement of NRAMP1 from *Arabidopsis thaliana* in iron transport. Biochem. J. **347,** 749-755.

Curie, C., Panaviene, Z., Loulergue, C., Dellaporta, S.L., Briat, J.F., Walker, E.L. (2001). Maize yellow stripe1 encodes a membrane protein directly involved in Fe(III) uptake. Nature **409,** 346-349.

Czarneckaverner, E., Yuan, C.X., Fox, P.C., Gurley, W.B. (1995). Isolation and Characterization of 6 Heat-Shock Transcription Factor cDNA Clones from Soybean. Plant Mol. Biol. **29,** 37-51.

Dahmani-Müller, H., Van Oort, F., B., G., M., B. (2000). Strategies of heavy metal uptake by three plant species growing near a metal smelter. Environ. Pollut. **109,** 231-238.

De Knecht, J.A., Van Baren, N., Ten Bookum, W.M., Wong Fong Sang, H.W., Koevoets, P.L.M., Schat, H., Verkleij, J.A.C. (1995). Synthesis and degradation of phytochelatins in Cd-sensitive and Cd-tolerant *Silene vulgaris*. Plant Sci. **106,** 9-18.

De Vos, C.H.R., Schat, H., Dewaal, M.A.M., Vooijs, R., Ernst, W.H.O. (1991). Increased Resistance to Copper-Induced Damage of the Root Cell Plasmalemma in Copper Tolerant *Silene cucubalus*. Physiol. Plant. **82,** 523-528.

Debouck, D. (1995). Differential display or differential dismay. Curr. Opin. Biol. **6,** 597-599.

Desikan, R., Clarke, A., Atherfold, P., Hancock, J.T., Neill, S.J. (1999b). Harpin induces mitogen-activated protein kinase activity during defence responses in *Arabidopsis thaliana* suspension cultures. Planta **210,** 97-103.

Desikan, R., Clarke, A., Hancock, J.T., Neill, S.J. (1999a). H_2O_2 activates a MAP kinase-like enzyme in *Arabidopsis thaliana* suspension cultures. J. Exp. Bot. **50,** 1863-1866.

Desikan, R., Hancock, J.T., Ichimura, K., Shinozaki, K., Neill, S.J. (2001). Harpin induces activation of the *Arabidopsis* mitogen-activated protein kinases AtMPK4 and AtMPK6. Plant Physiol. **126,** 1579-1587.

Desikan, R., S, A.H.-M., Hancock, J.T., Neill, S.J. (2001). Regulation of the *Arabidopsis* transcriptome by oxidative stress. Plant Physiol. **127,** 159-172.

Devetten, N.C., Ferl, R.J. (1995). Characterization of a Maize G-Box Binding-Factor That Is Induced by Hypoxia. Plant J. **7,** 589-601.

DeWald, D.B., Torabinejad, J., Jones, C.A., Shope, J.C., Cangelosi, A.R., Thompson, J.E., Prestwich, G.D., Hama, H. (2001). Rapid accumulation of phosphatidylinositol 4,5-bisphosphate and inositol 1,4,5-trisphosphate correlates with calcium mobilization in salt-stressed *Arabidopsis*. Plant Physiol. **126,** 759-769.

Diatchenko, L., Lau, Y.F., Campbell, A.P., Chenchik, A., Moqadam, F., Huang, B., Lukyanov, S., Gurskaya, N., Sverdlov, E.D., Siebert, P.D. (1996). Suppression subtractive hybridization: a method for generating differentially regulated or tissue-specific cDNA probes and libraries. Proc. Natl. Acad. Sci. USA **12,** 6025-6030.

Ditt, R.F., Nester, E.W., Comai, L. (2001). Plant gene expression response to *Agrobacterium tumefaciens*. Proc. Natl. Acad. Sci. USA **98,** 10954-10959.

Doyle, L.A., Yang, W., Abruzzo, L.V., Krogmann, T., Gao, Y., Rishi, A.K., Ross, D.D. (1998). A multidrug resistance transporter from human MCF-7 breast cancer cells. Proc. Natl. Acad. Sci. USA **95,** 15665-15670.

Dreesen, T.D., Johnson, D.H., Henikoff, S. (1988). The brown protein of *Drosophila melanogaster* is similar to the white protein and to components of active transport complexes. Mol. Cell Biol. **8,** 5206-5215.

Droillard, M., Boudsocq, M., Barbier-Brygoo, H., Lauriere, C. (2002). Different protein kinase families are activated by osmotic stresses in *Arabidopsis thaliana* cell suspensions. Involvement of the MAP kinases AtMPK3 and AtMPK6. FEBS Lett. **527,** 43-50.

Droillard, M.J., Thibivilliers, S., Cazale, A.C., Barbier-Brygoo, H., Lauriere, C. (2000). Protein kinases induced by osmotic stresses and elicitor molecules in tobacco cell suspensions: two crossroad MAP kinases and one osmoregulation-specific protein kinase. FEBS Lett. **474,** 217-222.

Dubos, C., Plomion, C. (2003). Identification of water-deficit responsive genes in maritime pine (*Pinus pinaster* Ait.) roots. Plant Mol. Biol. **51,** 249-262.

Durrant, W.E., Rowland, O., Piedras, P., Hammond-Kosack, K.E., Jones, J.D. (2000). cDNA-AFLP reveals a striking overlap in race-specific resistance and wound response gene expression profiles. Plant Cell **12,** 963-977.

Ebbs, S., Lau, I., Ahner, B., Kochian, L. (2002). Phytochelatin synthesis is not responsible for Cd tolerance in the Zn/Cd hyperaccumulator *Thlaspi caerulescens* (J. & C. Presl). Planta **214,** 635-640.

Eide, D., Broderius, M., Fett, J., Guerinot, M.L. (1996). A novel iron-regulated metal transporter from plants identified by functional expression in yeast. Proc. Natl. Acad. Sci. USA **93,** 5624-5628.

Elion, E.A. (1998). Routing MAP kinase cascades. Science **281,** 1625-1626.

Elion, E.A. (2000). Pheromone response, mating and cell biology. Curr. Opin. Microbiol. **3,** 573-581.

Elion, E.A. (2001). The Ste5p scaffold. J. Cell Sci. **114,** 3967-3978.

Enard, W., Khaitovich, P., Klose, J., Zollner, S., Heissig, F., Giavalisco, P., Nieselt-Struwe, K., Muchmore, E., Varki, A., Ravid, R., Doxiadis, G.M., Bontrop, R.E., Paabo, S. (2002). Intra- and interspecific variation in primate gene expression patterns. Science **296,** 340-343.

Ernst, W. (1974). Schwermetallvegetationen der Erde. (Stuttgart: G. Fischer Verlag).

Ernst, W.H.O., Joosse-van Damme, E.N.G. (1983). Umweltbelastungen durch Mineralstoffe - Biologische Effekte-. (VEB Gustav Fischer Verlag Jena).

Ewing, R.M., Kahla, A.B., Poirot, O., Lopez, F., Audic, S., Claverie, J.M. (1999). Large-scale statistical analysis of rice ESTs revealed correlated patterns of gene expression. Genome Res. **9,** 950-959.

Fairbanks, G., Steck, T.L., Wallach, D.F.H. (1971). Electrophoretic analysis of the major polypeptides of the human erythrocyte membrane. Biochemistry **10,** 2606-2616.

Falquet, L., Pagni, M., Bucher, P., Hulo, N., Sigrist, C.J., Hofmann, K., Bairoch, A. (2002). The PROSITE database, its status in 2002. Nucl. Acids Res. **30,** 235-238.

Fausto da Silva, J.J.R., Williams, R.J.P. (2001). The Biological Chemistry of the Elements. (Oxford University Press).

Felsenstein, J. (1993). PHYLIP (Phylogeny Inference Package) (Seattle: Depatment of Genetics University of Washington).

Fojtova, M., Kovarik, A. (2000). Genotoxic effect of cadmium is associated with apoptotic changes in tobacco cells. Plant Cell Environ. **23,** 531-537.

Fox, T.C., Guerinot, M.L. (1998). Molecular biology of cation transport in plants. Annu. Rev. Plant Physiol. Plant Mol. Biol. **49,** 669-696.

Fukazawa, J., Sakai, T., Ishida, S., Yamaguchi, I., Kamiya, Y., Takahashi, Y. (2000). REPRESSION OF SHOOT GROWTH, a bZIP transcriptional activator, regulates cell elongation by controlling the level of gibberellins. Plant Cell **12,** 901-915.

Fukuda, T., Kido, A., Kajino, K., Tsutsumi, M., Miyauchi, Y., Tsujiuchi, T., Konishi, Y., Hino, O. (1999). Cloning of Differentially Expressed Genes in Highly and Low Metastatic Rat Osteosarcmas by a Modified cDNA-AFLP Method. Biochem. and Biophys. Res. Commun. **261,** 35-40.

Gamas, P., Niebel, F.d.C., Lescure, N., Cullimore, J. (1996). Use of a subtractive hybridization approach to identify new *Medicago truncatula* genes induced during root nodule development. Mol. Plant Microbe Interact. **9,** 233-242.

Garcia-Hernandez, M., Murphy, A., Taiz, L. (1998). Metallothioneins 1 and 2 have distinct but overlapping expression patterns in *Arabidopsis*. Plant Physiol. **118,** 387-397.

Garcia-Olmedo, F., Molina, A., Alamillo, J.M., Rodriguez-Palenzuela, P. (1998). Plant defense peptides. Biopolymers **47,** 479-491.

Gerke, V., Moss, S.E. (2002). Annexins: from structure to function. Physiol. Rev. **82,** 331-371.

Ghassemian, M., Waner, D., Tchieu, J., Gribskov, M., Schroeder, J. I. (2001). An integrated *Arabidopsis* annotation database for Affymetrix Genechip data analysis, and tools for regulatory motif searches. Trends Plant Sci. **6**, 448-449.

Giedroc, D.P., Chen, X., Apuy, J.L. (2001). Metal response element (MRE)-binding transcription factor-1 (MTF-1): structure, function, and regulation. Antioxid. Redox Signal. **3,** 577-596.

Girke, T., Todd, J., Ruuska, S., White, J., Benning, C., Ohlrogge, J. (2000). Microarray analysis of developing *Arabidopsis* seeds. Plant Physiol. **124,** 1570-1581.

Glerum, D.M., Shtanko, A., Tzagoloff, A. (1996). Characterization of COX17, a yeast gene involved in copper metabolism and assembly of cytochrome oxidase. J. Biol. Chem. **271,** 14504-14509.

Granger, C.L., Cyr, R.J. (2001). Characterization of the yeast copper-inducible promoter system in *Arabidopsis thaliana*. Plant Cell Rep. **20,** 227-234.

Grill, E., Loffler, S., Winnacker, E.L., Zenk, M.H. (1989). Phytochelatins, the Heavy-Metal-Binding Peptides of Plants, Are Synthesized from Glutathione by a Specific Gamma-Glutamylcysteine Dipeptidyl Transpeptidase (Phytochelatin Synthase). Proc. Natl. Acad. Sci. USA **86,** 6838-6842.

Grill, E., Winnacker, E.L., Zenk, M.H. (1985). Phytochelatins - the Principal Heavy-Metal Complexing Peptides of Higher-Plants. Science **230,** 674-676.

Grotz, N., Fox, T., Connolly, E., Park, W., Guerinot, M.L., Eide, D. (1998). Identification of a family of zinc transporter genes from *Arabidopsis* that respond to zinc deficiency. Proc. Natl. Acad. Sci. USA **95,** 7220-7224.

Guerinot, M.L. (2000). The ZIP family of metal transporters. Biochim. Biophys. Acta Biomembr. **1465,** 190-198.

Ha, S.B., Smith, A.P., Howden, R., Dietrich, W.M., Bugg, S., O'Connell, M.J., Goldsbrough, P.B., Cobbett, C.S. (1999). Phytochelatin synthase genes from *Arabidopsis* and the yeast *Schizosaccharomyces pombe*. Plant Cell **11,** 1153-1163.

Haag, E., Raman, V. (1994). Effects of primer choice and source of Taq DNA polymerase on the banding patterns of differential display RT-PCR. Biotechniques **2,** 226-228.

Habu, Y., Fukada-Tanaka, S., Hisatomi, Y., Iida, S. (1997). Amplified restriction fragment length polymorphism-based mRNA fingerprinting using a single restriction enzyme that recognizes a 4-bp sequence. Biochem. Biophys. Res. Commun. **234,** 516-521.

Hamer, D.H. (1986). Metallothionein. Annu. Rev. Biochem. **55,** 913-951.

Hamer, D.H., Thiele, D.J., Lemontt, J.E. (1985). Function and autoregulation of yeast copperthionein. Science **228,** 685-690.

Harada, E., Yamaguchi, Y., Koizumi, N., Sano, H. (2002). Cadmium stress induces production of thiol compounds and transcripts for enzymes involved in sulfur assimilation pathways in *Arabidopsis*. J. Plant Physiol. **159,** 445-448.

Harres, H.P. (1999). Schwermetallgehalte in Böden, Pflanzen, Fließgewässern und Sedimenten unterschiedlicher belasteter Regionen Südsardiniens. In Mathematisch-Naturwissenschaftlich-Technischen Fakultät (Halle/S.: Martin-Luther-Universität Halle-Wittenberg).

Hasegawa, M., Bressan, R., Pardo, J.M. (2000). The dawn of plant salt tolerance genetics. Trends Plant Sci. **5,** 317-319.

Heider, H., Boscheinen, O., Scharf, K.D. (1998). A heat-stress pulse inactivates a 50 kDa myelin basic protein kinase in tomato. Bot. Acta **111,** 398-401.

Heiss, S., Schafer, H.J., Haag-Kerwer, A., Rausch, T. (1999). Cloning sulfur assimilation genes of *Brassica juncea* L.: cadmium differentially affects the expression of a putative low-affinity sulfate transporter and isoforms of ATP sulfurylase and APS reductase. Plant Mol. Biol. **39,** 847-857.

Hell, R., Stephan, U.W. (2003). Iron uptake, trafficking and homeostasis in plants. Planta **216,** 541-551.

Henriques, R., Jasik, J., Klein, M., Martinoia, E., Feller, U., Schell, J., Pais, M.S., Koncz, C. (2002). Knock-out of *Arabidopsis* metal transporter gene *IRT1* results in iron deficiency accompanied by cell differentiation defects. Plant Mol. Biol. **50,** 587-597.

Himelblau, E., Mira, H., Lin, S.J., Culotta, V.C., Penarrubia, L., Amasino, R.M. (1998). Identification of a functional homolog of the yeast copper homeostasis gene ATX1 from *Arabidopsis*. Plant Physiol. **117,** 1227-1234.

Hirayama, T., Kieber, J.J., Hirayama, N., Kogan, M., Guzman, P., Nourizadeh, S., Alonso, J.M., Dailey, W.P., Dancis, A., Ecker, J.R. (1999). Responsive-to-antagonist1, a Menkes/Wilson disease-related copper transporter, is required for ethylene signaling in *Arabidopsis*. Cell **97,** 383-393.

Hirayama, T., Ohto, C., Mizoguchi, T., Shinozaki, K. (1995). A gene encoding a phosphatidylinositol-specific phospholipase C is induced by dehydration and salt stress in *Arabidopsis thaliana*. Proc. Natl. Acad. Sci. USA **92,** 3903-3907.

Hirota, K., Murata, M., Sachi, Y., Nakamura, H., Takeuchi, J., Mori, K., Yodoi, J. (1999). Distinct roles of thioredoxin in the cytoplasm and in the nucleus. A two-step mechanism of redox regulation of transcription factor NF-kappaB. J. Biol. Chem. **274,** 27891-27897.

Hirschi, K.D. (1999). Expression of *Arabidopsis* CAX1 in tobacco: Altered calcium homeostasis and increased stress sensitivity. Plant Cell **11,** 2113-2122.

Hoffmann, R., Seidl, T., Dugas, M. (2002). Profound effect of normalization on detection of differentially expressed genes in oligonucleotide microarray data analysis. Genome Biol. **3,** 1-11.

Hofmann, A., Proust, J., Dorowski, A., Schantz, R., Huber, R. (2000). Annexin 24 from *Capsicum annuum*. X-ray structure and biochemical characterization. J. Biol. Chem. **275,** 8072-8082.

Holm, M., Ma, L.G., Qu, L.J., Deng, X.W. (2002). Two interacting bZIP proteins are direct targets of COP1-mediated control of light-dependent gene expression in *Arabidopsis*. Genes Dev. **16,** 1247-1259.

Horvath, D.M., Chua, N.-H. (1996). Identification of an immediate-early salicylic acid-inducible tobacco gene and characterization of induction by other compounds. Plant Mol. Biol. **31,** 1061-1072.

Howden, R., Goldsbrough, P.B., Andersen, C.R., Cobbett, C.S. (1995). Cadmium-sensitive, cad1 mutants of *Arabidopsis thaliana* are phytochelatin deficient. Plant Physiol. **107,** 1059-1066.

Huala, E., Dickerman, A.W., Garcia-Hernandez, M., Weems, D., Reiser, L., LaFond, F., Hanley, D., Kiphart, D., Zhuang, M., Huang, W., Mueller, L.A., Bhattacharyya, D., Bhaya, D., Sobral, B.W., Beavis, W., Meinke, D.W., Town, C.D., Somerville, C., Rhee, S.Y. (2001). The Arabidopsis Information Resource (TAIR): a comprehensive database and web-based information retrieval, analysis, and visualization system for a model plant. Nucl. Acids Res. **29,** 102-105.

Ichimura, K., Mizoguchi, T., Yoshida, R., Yuasa, T., Shinozaki, K. (2000). Various abiotic stresses rapidly activate *Arabidopsis* MAP kinases ATMPK4 and ATMPK6. Plant J. **24,** 655-665.

Ichimura, K., Shinozaki, K., Tena, G., Sheen, J., Henry, Y., Champion, A., Kreis, M., Zhang, S., Hirt, H., Wilson, C., Heberle-Bors, E., Ellis, B.E., Morris, P.C., Innes, R.W., Ecker, J.E., Scheel, D., Klessig, D.F., Y., M., Mundy, J., Ohashi, Y., Walker, J.C. (2002). Mitogen-activated protein kinase cascades in plants: a new nomenclature. Trends Plant Sci. **7,** 301-308.

Izawa, T., Foster, R., Chua, N.H. (1993). Plant bZIP Protein-DNA Binding-Specificity. J. Mol. Biol. **230,** 1131-1144.

Jakoby, M., Weisshaar, B., Droge-Laser, W., Vicente-Carbajosa, J., Tiedemann, J., Kroj, T., Parcy, F. (2002). bZIP transcription factors in *Arabidopsis*. Trends Plant Sci. **7,** 106-111.

Johnson, L. N., O'Reilly, M. (1996). Control by phosphorylation. Curr. Opin. Struct. Biol. **6**, 762-769

Jonak, C., Kiegerl, S., Ligterink, W., Barker, P.J., Huskisson, N.S., Hirt, H. (1996). Stress signaling in plants: a mitogen-activated protein kinase pathway is activated by cold and drought. Proc. Natl. Acad. Sci. USA **93,** 11274-11279.

Jonak, C., Okresz, L., Bogre, L., Hirt, H. (2002). Complexity, cross talk and integration of plant MAP kinase signalling. Curr. Opin. Plant Biol. **5,** 415-424.

Kadyrzhanova, D.K., Vlachonasios, K.E., Ververidis, P., Dilley, D.R. (1998). Molecular cloning of a novel heat induced/ chilling tolerance related cDNA in tomato fruit by use of mRNA differential display. Plant Mol. Biol. **36,** 885-895.

Kamizono, A., Nishizawa, M., Teranishi, Y., Murata, K., Kimura, A. (1989). Identification of a Gene Conferring Resistance to Zinc and Cadmium Ions in the Yeast *Saccharomyces cerevisiae*. Mol. Gen. Genet. **219,** 161-167.

Kampfenkel, K., Kushnir, S., Babiychuk, E., Inze, D., Vanmontagu, M. (1995). Molecular Characterization of a Putative *Arabidopsis thaliana* Copper Transporter and Its Yeast Homolog. J. Biol. Chem. **270,** 28479-28486.

Kang, J.Y., Choi, H.I., Im, M.Y., Kim, S.Y. (2002). *Arabidopsis* basic leucine zipper proteins that mediate stress-responsive abscisic acid signaling. Plant Cell **14,** 343-357.

Kasuga, M., Liu, Q., Miura, S., Yamaguchi-Shinozaki, K., Shinozaki, K. (1999). Improving plant drought, salt, and freezing tolerance by gene transfer of a single stress-inducible transcription factor. Nat. Biotechnol.**17,** 287-291.

Kemmerling, B. (2001). Identifizierung, Charakterisierung systemisch responsiver Gene der Kartoffel (*Solanum tuberosum* L.) nach Inokulation mit dem nichtpathogenen Bakterium *Pseudomonas syringae pv. maculicola*. (Halle/S.: Martin-Luther-Universität).

Kerk, D., Bulgrien, J., Smith, D.W., Barsam, B., Veretnik, S., Gribskov, M. (2002). The complement of protein phosphatase catalytic subunits encoded in the genome of *Arabidopsis*. Plant Physiol. **129,** 908-925.

Kerr, I.D. (2002). Structure and association of ATP-binding cassette transporter nucleotide-binding domains. Biochim. Biophys. Acta **1561,** 47-64.

Kieber, J.J., Rothenberg, M., Roman, G., Feldmann, K.A., Ecker, J.R. (1993). Ctr1, a Negative Regulator of the Ethylene Response Pathway in *Arabidopsis*, Encodes a Member of the Raf Family of Protein- Kinases. Cell **72,** 427-441.

Kiegerl, S., Cardinale, F., Siligan, C., Gross, A., Baudouin, E., Liwosz, A., Eklof, S., Till, S., Bogre, L., Hirt, H., Meskiene, I. (2000). SIMKK, a mitogen-activated protein kinase (MAPK) kinase, is a specific activator of the salt stress-induced MAPK, SIMK. Plant Cell **12,** 2247-2258.

Kim, H.S., Delaney, T.P. (2002). Over-expression of *TGA5*, which encodes a bZIP transcription factor that interacts with NIM1/NPR1, confers SAR-independent resistance in *Arabidopsis thaliana* to *Peronospora parasitica*. Plant J. **32,** 151-163.

Kim, J.C., Lee, S.H., Cheong, Y.H., Yoo, C.M., Lee, S.I., Chun, H.J., Yun, D.J., Hong, J.C., Lee, S.Y., Lim, C.O., Cho, M.J. (2001). A novel cold-inducible zinc finger protein from soybean, SCOF-1, enhances cold tolerance in transgenic plants. Plant J. **25,** 247-259.

Kistner, C. (1999). Isolierung und Charakterisierung eines nach Infektion mit *Pseudomonas syringae pv. maculicola* systemisch responsiven Gens von *Solanum tuberosum* L. (Halle/S.: Martin-Luther-Universität).

Knetsch, M.L.W., Wang, M., Snaar-Jagalska, B.E., Heimovaara-Dijkstra, S. (1996). Abscisic acid induces mitogen-activated protein kinase activation in barley aleurone protoplasts. Plant Cell **8,** 1061-1067.

Knight, H. (2000). Calcium signaling during abiotic stress in plants. Int. Rev. Cytol. **195,** 269-324.

Koch, M., Bishop, J., Mitchell-Olds, T. (1999). Molecular systematics and evolution of *Arabidopsis* and *Arabis*. Plant Biol. **1,** 529-537.

Koch, M.A., Haubold, B., Mitchell-Olds, T. (2000). Comparative Evolutionary Analysis of Chalcone Synthase ans Alcohol Dehydrogenase Loci in *Arabidopsis*, *Arabis*, and Related Genera (*Brassicaceae*). Mol. Biol. Evol. **17,** 1483-1498.

Koch, M.A., Weisshaar, B., Kroymann, J., Haubold, B., Mitchell-Olds, T. (2001). Comparative Genomics and Regulatory Evolution: Conservation and Function of the *Chs* and *Apetala3* Promoters. Mol. Biol. Evol. **18,** 1882-1891.

Koga, M., Zwaal, R., Guan, K.L., Avery, L., Ohshima, Y. (2000). A *Caenorhabditis elegans* MAP kinase kinase, MEK-1, is involved in stress responses. Embo J. **19,** 5148-5156.

Koncz, C., Schell, J. (1986). The promotor of TL-DNA gene 5 controls the tissue-specific expression of chimeric genes carried by a novel type of *Agrobacterium* binary vector. Mol. Gen. Genet. **204,** 383-396.

Korshunova, Y.O., Eide, D., Clark, W.G., Guerinot, M.L., Pakrasi, H.B. (1999). The IRT1 protein from *Arabidopsis thaliana* is a metal transporter with a broad substrate range. Plant Mol. Biol. **40,** 37-44.

Kovtun, Y., Chiu, W.L., Tena, G., Sheen, J. (2000). Functional analysis of oxidative stress-activated mitogen-activated protein kinase cascade in plants. Proc. Natl. Acad. Sci. USA **97,** 2940-2945.

Krämer, U., Cotter Howells, J.D., Charnock, J.M., Baker, A.J.M., Smith, J.A.C. (1996). Free histidine as a metal chelator in plants that accumulate nickel. Nature **379,** 635-638.

Krämer, U., Pickering, I.J., Prince, R.C., Raskin, I., Salt, D.E. (2000). Subcellular localization and speciation of nickel in hyperaccumulator and non-accumulator *Thlaspi* species. Plant Physiol. **122,** 1343-1353.

Kreps, J.A., Wu, Y., Chang, H.S., Zhu, T., Wang, X., Harper, J.F. (2002). Transcriptome changes for *Arabidopsis* in response to salt, osmotic, and cold stress. Plant Physiol. **130,** 2129-2141.

Kroj, T., Rudd, J.J., Nürnberger, T., Gäbler, Y., Lee, J., Scheel, D. (2003). Mitogen-activated Protein Kinases Play an Essential Role in Oxidative Burst-independent Expression of Pathogenesis-related Genes in Parsley. J. Biol. Chem. **278,** 2256-2264.

Krupa, Z. (1988). Cadmium-Induced Changes in the Composition and Structure of the Light-Harvesting Chlorophyll a/b Protein Complex-II in Radish Cotyledons. Physiol. Plant. **73,** 518-524.

Krysan, P.J., Young, J.C., Sussman, M.R. (1999). T-DNA as an Insertional Mutagen in *Arabidopsis*. Plant Cell **11,** 2283-2290.

Küpper, H., Lombi, E., Zhao, F.J., McGrath, S.P. (2000). Cellular compartmentation of cadmium and zinc in relation to other elements in the hyperaccumulator *Arabidopsis halleri*. Planta **212,** 75-84.

Kusano, T., Berberich, T., Harada, M., Suzuki, N., Sugawara, K. (1995). A Maize DNA-Binding Factor with a bZIP Motif Is Induced by Low-Temperature. Mol. Gen. Genet. **248,** 507-517.

Kyte, J., Dolittle, F.R. (1982). A simple method for displaying the hydropathic character of a protein. J. Mol. Biol. **157,** 693-698.

Labbe, S., Zhu, Z., Thiele, D.J. (1997). Copper-specific transcriptional repression of yeast genes encoding critical components in the copper transport pathway. J. Biol. Chem. **272,** 15951-15958.

Laity, J.H., Lee, B.M., Wright, P.E. (2001). Zinc finger proteins: new insights into structural and functional diversity. Curr. Opin. Struct. Biol. **11,** 39-46.

Lane, B., Kajioka, R., Kennedy, T. (1987). The Wheat-Germ-Ec Protein Is a Zinc-Containing Metallothionein. Biochem. Cell Biol. **65,** 1001-1005.

Lane, T.W., Morel, F.M.M. (2000). A biological function for cadmium in marine diatoms. Proc. Natl. Acad. Sci. USA **97,** 4627-4631.

Larsson, E.H., Bornman, J.F., Asp, H. (1998). Influence of UV-B radiation and Cd^{2+} on chlorophyll fluorescence, growth and nutrient content in *Brassica napus*. J. Exp. Bot. **49,** 1031-1039.

Lasat, M.M., Baker, A., Kochian, L.V. (1996). Physiological Characterization of Root Zn^{2+} Absorption and Translocation to Shoots in Zn Hyperaccumulator and Nonaccumulator Species of *Thlaspi*. Plant Physiol. **112,** 1715-1722.

Lasat, M.M., Baker, A.J., Kochian, L.V. (1998). Altered Zn compartmentation in the root symplasm and stimulated Zn absorption into the leaf as mechanisms involved in Zn hyperaccumulation in *Thlaspi caerulescens*. Plant Physiol. **118,** 875-883.

Lee, B.N., Elion, E.A. (1999). The MAPKKK Ste11 regulates vegetative growth through a kinase cascade of shared signaling components. Proc. Natl. Acad. Sci. USA **96,** 12679-12684.

Lee, J., Klessig, D.F., Nürnberger, T. (2001). A harpin binding site in tobacco plasma membranes mediates activation of the pathogenesis-related gene HIN1 independent of extracellular calcium but dependent on mitogen-activated protein kinase activity. Plant Cell **13,** 1079-1093.

Lee, S., Korban, S.S. (2002). Transcriptional regulation of *Arabidopsis thaliana* phytochelatin synthase (*AtPCS1*) by cadmium during early stages of plant development. Planta **215,** 689-693.

Lee, S.J., Lee, M.Y., Yi, S.Y., Sang, K.H., Choi, S.H., Her, N.H., Choi, D., Min, B.W., Yang, S.G., Harn, C.H. (2002). PPI1: A novel pathogen-induced basic region-leucine zipper (bZIP) transcription factor from pepper. Mol. Plant Microbe Interac. **15,** 540-548.

Lemaire, S., Keryer, E., Stein, M., Schepens, I.I., Issakidis-Bourguet, E., Grard-Hirne, C., Miginiac-Maslow, M., Jacquot, J.P. (1999). Heavy-metal regulation of thioredoxin gene expression in *Chlamydomonas reinhardtii*. Plant Physiol. **120,** 773-778.

Lescot, M., Dehais, P., Thijs, G., Marchal, K., Moreau, Y., Van de Peer, Y., Rouze, P., Rombauts, S. (2002). PlantCARE, a database of plant cis-acting regulatory elements and a portal to tools for in silico analysis of promoter sequences. Nucl. Acids Res. **30,** 325-327.

Leung, J., Bouvierdurand, M., Morris, P.C., Guerrier, D., Chefdor, F., Giraudat, J. (1994). *Arabidopsis* ABA Response Gene ABI1 - Features of a Calcium- Modulated Protein Phosphatase. Science **264,** 1448-1452.

Leung, J., Merlot, S., Giraudat, J. (1997). The *Arabidopsis* ABSCISIC ACID-INSENSITIVE2 (ABI2) and ABI1 Genes Encode Homologous Protein Phosphatases 2C Involved in Abscisic Acid Signal Transduction. Plant Cell **9,** 759-771.

Li, H.Y., Gray, J.E. (1997). Pollination-enhanced expression of a receptor-like protein kinase related gene in tabacco styles. Plant Mol. Biol. **33,** 653-665.

Li, L.G., He, Z.Y., Pandey, G.K., Tsuchiya, T., Luan, S. (2002). Functional cloning and characterization of a plant efflux carrier for multidrug and heavy metal detoxification. J. Biol. Chem. **277,** 5360-5368.

Li, L.G., Tutone, A.F., Drummond, R.S.M., Gardner, R.C., Luan, S. (2001). A novel family of magnesium transport genes in *Arabidopsis*. Plant Cell **13,** 2761-2775.

Li, L.T., Kaplan, J. (1998). Defects in the yeast high affinity iron transport system result in increased metal sensitivity because of the increased expression of transporters with a broad transition metal specificity. J. Biol. Chem. **273,** 22181-22187.

Li, Z.S., Lu, Y.P., Zhen, R.G., Szczypka, M., Thiele, D.J., Rea, P.A. (1997). A new pathway for vacuolar cadmium sequestration in *Saccharomyces cerevisiae*: YCF1-catalyzed transport of bis(glutathionato)cadmium. Proc. Natl. Acad. Sci. USA **94,** 42-47.

Liang, P., Pardee, A.B. (1992). Differential display of eucaryotic messenger RNA by means of the polymerase chain reaction. Science **257,** 967-971.

Lichtenthaler, H.K., Wellburn, A.R. (1983). Determination of the total carotinoids and chlorophylls a and b of leaf extracts in different solvents. Biochem. Soc. Transac. **603,** 591-592.

Lin, S.J., Culotta, V.C. (1995). The *Atx1* Gene of *Saccharomyces cerevisiae* Encodes a Small Metal Homeostasis Factor That Protects Cells against Reactive Oxygen- Toxicity. Proc. Natl. Acad. Sci. USA **92,** 3784-3788.

Liu, L.S., White, M.J., MacRae, T.H. (1999). Transcription factors and their genes in higher plants - Functional domains, evolution and regulation. Eur. J. Biochem. **262,** 247-257.

Liu, Q., Kasuga, M., Sakuma, Y., Abe, H., Miura, S., Yamaguchi-Shinozaki, K., Shinozaki, K. (1998). Two transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought- and low-temperature-responsive gene expression, respectively, in *Arabidopsis*. Plant Cell **10,** 1391-1406.

Lopez-Molina, L., Chua, N.H. (2000). A null mutation in a bZIP factor confers ABA-insensitivity in *Arabidopsis thaliana*. Plant Cell Physiol. **41,** 541-547.

Luan, S. (2003). Protein Phosphatases in Plants. Annu. Rev. Plant Biol. **54**, 63-92

Lyons, T.J., Gasch, A.P., Gaither, L.A., Botstein, D., Brown, P.O., Eide, D.J. (2000). Genome-wide characterization of the Zap1p zinc-responsive regulon in yeast. Proc. Natl. Acad. Sci. USA **97,** 7957-7962.

Macnair, M.R. (2002). Within and between population genetic variation for zinc accumulation in *Arabidopsis halleri*. New Phytol. **155,** 59-66.

Macnair, M.R., Bert, V., Huitson, S.B., Saumitou-Laprade, P., Petit, D. (1999). Zinc tolerance and hyperaccumulation are genetically independent characters. Proc. R. Soc. Lond. B **266,** 2175-2179.

Marschner, H. (1995). Mineral Nutrition of Higher Plants. (London: Academic Press).

Marshall, C.J. (1995). Specificity of receptor tyrosine kinase signaling: transient versus sustained extracellular signal-regulated kinase activation. Cell **80,** 179-185.

Martinoia, E., Klein, M., Geisler, M., Bovet, L., Forestier, C., Kolukisaoglu, U., Muller-Rober, B., Schulz, B. (2002). Multifunctionality of plant ABC transporters-more than just detoxifiers. Planta **214,** 345-355.

Masters, B.A., Kelly, E.J., Quaife, C.J., Brinster, R.L., Palmiter, R.D. (1994). Targeted disruption of metallothionein I and II genes increases sensitivity to cadmium. Proc. Natl. Acad. Sci. USA **91,** 584-588.

Matsumura, H., Nirasawa, S., Terauchi, R. (1999). Transcript profiling in rice (*Oryza sativa*, L.) seedlings using serial analysis of gene expression (SAGE). Plant J. **20,** 719-726.

Matz, M.V., Lukyanov, S.A. (1998). Differential strategies of differential display: areas of application. Nulc. Acids Res. **26,** 5537-5543.

Mench, M., Morel, J.L., Guckert, A., Guillet, B. (1988). Metal-Binding with Root Exudates of Low-Molecular Weight. J. Soil. Sci. **39,** 521-527.

Merlot, S., Gosti, F., Guerrier, D., Vavasseur, A., Giraudat, J. (2001). The ABI1 and ABI2 protein phosphatases 2C act in a negative feedback regulatory loop of the abscisic acid signalling pathway. Plant J. **25,** 295-303.

Meskiene, I., Bögre, L., Glaser, W., Balog, J., Brandstötter, M., Zwerger, K., Ammerer, G., Hirt, H. (1998). MP2C, a plant protein phosphatase 2C, functions as a negative regulator of mitogen-activated protein kinase pathways in yeast and plants. Proc. Natl. Acad. Sci. USA **95,** 1938-1943.

Mikami, K., Katsura, M., Ito, T., Okada, K., Shimura, Y., Iwabuchi, M. (1995). Developmental and Tissue-Specific Regulation of the Gene for the Wheat Basic/Leucine Zipper Protein Hbp-1a(17) in Transgenic *Arabidopsis* Plants. Mol. Gen. Genet. **248,** 573-582.

Mira, H., Martinez-Garcia, F., Penarrubia, L. (2001). Evidence for the plant-specific intercellular transport of the *Arabidopsis* copper chaperone CCH. Plant J. **25,** 521-528.

Miyashita, N.T., Innan, H., Terauchi, R. (1996). Intra- and Interspecific Variation of the Alcohol Dehydrogenase Locus Region in Wild Plants *Arabis gemmifera* and *Arabidopsis thaliana*. Mol. Biol. Evol. **13,** 433-436.

Mizoguchi, T., Irie, K., Hirayama, T., Hayashida, N., Yamaguchi-Shinozaki, K., Matsumoto, K., Shinozaki, K. (1996). A gene encoding a mitogen-activated protein kinase kinase kinase is induced simultaneously with genes for a mitogen-activated protein kinase and an S6 ribosomal protein kinase by touch, cold, and water stress in *Arabidopsis thaliana*. Proc. Natl. Acad. Sci. USA **93,** 765-769.

Mohr, K. (1978). Geologie und Minerallagerstätten des Harzes. (Stuttgart: E. Schweizerbart'sche Verlagsbuchhandlung).

Mouaheb, N., Thomas, D., Verdoucq, L., Monfort, P., Meyer, Y. (1998). In vivo functional discrimination between plant thioredoxins by heterologous expression in the yeast *Saccharomyces cerevisiae*. Proc. Natl. Acad. Sci. USA **95,** 3312-3317.

Munnik, T., Irvine, R.F., Musgrave, A. (1998). Phospholipid signalling in plants. Biochim. Biophys. Acta **1389,** 222-272.

Munnik, T., Ligterink, W., Meskiene, I.I., Calderini, O., Beyerly, J., Musgrave, A., Hirt, H. (1999). Distinct osmo-sensing protein kinase pathways are involved in signalling moderate and severe hyper-osmotic stress. Plant J. **20,** 381-388.

Murata, N., Los, D.A. (1997). Membrane Fluidity and Temperature Perception. Plant Physiol. **115,** 875-879.

Murphy, A., Taiz, L. (1995). Comparison of Metallothionein Gene-Expression and Nonprotein Thiols in 10 *Arabidopsis* Ecotypes - Correlation with Copper Tolerance. Plant Physiol. **109,** 945-954.

Murphy, A., Taiz, L. (1997). Correlation between potassium efflux and copper sensitivity in 10 *Arabidopsis* ecotypes. New Phytol. **136,** 211-222.

Nakai, K., Horton, P. (1999). PSORT: A program for detecting sorting signals in proteins and predicting their subcellular localization. Trends Biochem. Sci. **24,** 34-36.

Nakashima, K., Shinwari, Z.K., Sakuma, Y., Seki, M., Miura, S., Shinozaki, K., Yamaguchi-Shinozaki, K. (2000). Organization and expression of two *Arabidopsis* DREB2 genes encoding DRE-binding proteins involved in dehydration- and high-salinity-responsive gene expression. Plant Mol. Biol. **42,** 657-665.

Negishi, T., Nakanishi, H., Yazaki, J., Kishimoto, N., Fujii, F., Shimbo, K., Yamamoto, K., Sakata, K., Sasaki, T., Kikuchi, S., Mori, S., Nishizawa, N.K. (2002). cDNA microarray analysis of gene expression during Fe-deficiency stress in barley suggests that polar transport of vesicles is implicated in phytosiderophore secretion in Fe-deficient barley roots. Plant J. **30,** 83-94.

Neill, S.J., Desikan, R., Clarke, A., Hurst, R.D., Hancock, J.T. (2002). Hydrogen peroxide and nitric oxide as signalling molecules in plants. J. Exp. Bot. **53,** 1237-1247.

Neumann, D., Lichtenberger, O., Gunther, D., Tschiersch, K., Nover, L. (1994). Heat-Shock Proteins Induce Heavy-Metal Tolerance in Higher- Plants. Planta **194,** 360-367.

Nies, D.H. (1999). Microbial heavy-metal resistance. Appl. Microbiol. Biotechnol. **51,** 730-750.

Niggeweg, R., Thurow, C., Kegler, C., Gatz, C. (2000). Tobacco transcription factor TGA2.2 is the main component of as-1-binding factor ASF-1 and is involved in salicylic acid- and auxin-inducible expression of as-1-containing target promoters. J. Biol. Chem. **275,** 19897-19905.

Noctor, G., Gomez, L., Vanacker, H., Foyer, C.H. (2002). Interactions between biosynthesis, compartmentation and transport in the control of glutathione homeostasis and signalling. J. Exp. Bot. **53,** 1283-1304.

Noel, L.F., Thieme, D., Nennstiel, D., Bonas, U. (2001). cDNA-AFLP analysis unravels a genome-wide hrpG-regulon in the plant pathogen *Xanthomonas campestris* pv. *vesicatoria*. Mol. Microbiol. **41,** 1271-1281.

Nover, L., Bharti, K., Doring, P., Mishra, S.K., Ganguli, A., Scharf, K.D. (2001). *Arabidopsis* and the heat stress transcription factor world: how many heat stress transcription factors do we need? Cell Stress Chaperones **6,** 177-189.

Nuhse, T.S., Peck, S.C., Hirt, H., Boller, T. (2000). Microbial elicitors induce activation and dual phosphorylation of the *Arabidopsis thaliana* MAPK 6. J. Biol. Chem. **275,** 7521-7526.

O´Kane, S.L., Al-Shehbaz, I.A. (1997). A Synopsis of *Arabidopsis* (Brassicaceae). Novon **7**, 323-327.

Olmos, E., Martinez-Solano, J.R., Piqueras, A., Hellin, E. (2003). Early steps in the oxidative burst induced by cadmium in cultured tobacco cells (BY-2 line). J. Exp. Bot. **54,** 291-301.

O'Rourke, S.M., Herskowitz, I. (1998). The Hog1 MAPK prevents cross talk between the HOG and pheromone response MAPK pathways in *Saccharomyces cerevisiae*. Genes Dev. **12,** 2874-2886.

Ortiz, D.F., Kreppel, L., Speiser, D.M., Scheel, G., McDonald, G., Ow, D.W. (1992). Heavy metal tolerance in the fission yeast requires an ATP-binding cassette-type vacuolar membrane transporter. Embo J. **11,** 3491-3499.

Ortiz, D.F., Ruscitti, T., McCue, K.T., Ow, D.W. (1995). Transport of Metal-binding Peptides by HMT1, A Fission Yeats ABC-type Vacuolar Membrane Protein. J. Biol. Chem. **270,** 4721-4728.

Ouariti, O., Boussama, N., Zarrouk, M., Cherif, A., Ghorbal, M.H. (1997). Cadmium- and copper-induced changes in tomato membrane lipids. Phytochemistry **45,** 1343-1350.

Outten, C.E., O'Halloran, T.V. (2001). Femtomolar sensitivity of metalloregulatory proteins controlling zinc homeostasis. Science **292,** 2488-2492.

Oyama, T., Shimura, Y., Okada, K. (1997). The *Arabidopsis HY5* gene encodes a bZIP protein that regulates stimulus-induced development of root and hypocotyl. Genes Dev. **11,** 2983-2995.

Pawson, T. (1995). Protein modules and signalling networks. Nature **373**, 573-580.

Pence, N.S., Larsen, P.B., Ebbs, S.D., Letham, D.L., Lasat, M.M., Garvin, D.F., Eide, D., Kochian, L.V. (2000). The molecular physiology of heavy metal transport in the Zn/Cd hyperaccumulator *Thlaspi caerulescens*. Proc. Natl. Acad. Sci. USA **97,** 4956-4960.

Perfus-Barbeoch, L., Leonhardt, N., Vavasseur, A., Forestier, C. (2002). Heavy metal toxicity: cadmium permeates through calcium channels and disturbs the plant water status. Plant J. **32,** 539-548.

Persans, M.W., Yan, X.G., Patnoe, J., Krämer, U., Salt, D.E. (1999). Molecular dissection of the role of histidine in nickel hyperaccumulation in *Thlaspi goesingense* (Halacsy). Plant Physiol. **121,** 1117-1126.

Petersen, M., Brodersen, P., Naested, H., Andreasson, E., Lindhart, U., Johansen, B., Nielsen, H.B., Lacy, M., Austin, M.J., Parker, J.E., Sharma, S.B., Klessig, D.F., Martienssen, R., Mattsson, O., Jensen, A.B., Mundy, J. (2000). *Arabidopsis* map kinase 4 negatively regulates systemic acquired resistance. Cell **103,** 1111-1120.

Petters, J. (2001). Isolierung und Charakterisierung pathogen- und stress-induzierter Gene von Kartoffel *Solanum tuberosum* L. (Halle/S.: Martin-Luther-Universität).

Petters, J., Göbel, C., Scheel, D., Rosahl, S. (2002). A Pathogen-Responsive cDNA from Potato Encodes a Protein with Homology to a Phosphate Starvation-Induced Phosphatase. Plant Cell Physiol. **43,** 1049-1053.

Pich, A., Scholz, G. (1996). Translocation of copper and other micronutrients in tomato plants (*Lycopersicon esculentum* Mill): Nicotianamine-stimulated copper transport in the xylem. J. Exp. Bot. **47,** 41-47.

Plieth, C., Hansen, U.P., Knight, H., Knight, M.R. (1999). Temperature sensing by plants: the primary characteristics of signal perception and calcium response. Plant J. **18,** 491-497.

Qin, L., Overmars, H., Helder, J., Popeijus, H., van der Voort, J.R., Groenink, W., van Koert, P., Schots, A., Bakker, J., Smant, G. (2000). An Efficient cDNA-AFLP-Based Strategy for the Identification of Putative Pathogenicity Factors form Potato Cyst Nematode *Globodera rostochiensis*. Mol. Plant Microbe Interact. **13**, 830-836.

Rae, T.D., Schmidt, P.J., Pufahl, R.A., Culotta, V.C., O'Halloran, T.V. (1999). Undetectable intracellular free copper: The requirement of a copper chaperone for superoxide dismutase. Science **284,** 805-808.

Rauser, W.E. (1999). Structure and function of metal chelators produced by plants - The case for organic acids, amino acids, phytin, and metallothioneins. Cell Biochem. Biophys. **31,** 19-48.

Rigaut, G., Shevchenko, A., Rutz, B., Wilm, M., Mann, M., Séraphin, B. (1999). A generic proteinpurification method for protein complex characterization and proteome exploration. Nat. Biotechnol. **17,** 1030-1032.

Roberts, R.L., Fink, G.R. (1994). Elements of a single MAP kinase cascade in *Saccharomyces cerevisiae* mediate two developmental programs in the same cell type: mating and invasive growth. Genes Dev. **8,** 2974-2985.

Robinson, N.J., Wilson, J.R., Turner, J.S. (1996). Expression of the type 2 metallothionein-like gene MT2 from *Arabidopsis thaliana* in Zn^{2+}-metallothionein-deficient *Synechococcus* PCC 7942: Putative role for MT2 in Zn^{2+} metabolism. Plant Mol. Biol. **30,** 1169-1179.

Rodriguez, F.I., Esch, J.J., Hall, A.E., Binder, B.M., Schaller, G.E., Bleecker, A.B. (1999). A copper cofactor for the ethylene receptor ETR1 from *Arabidopsis*. Science **283,** 996-998.

Rodriguez, P.L. (1998). Protein phosphatases 2C (PP2C) function in higher plants. Plant Mol. Biol. **38,** 919-927.

Romeis, T., Piedras, P., Zhang, S., Klessig, D.F., Hirt, H., Jones, J.D. (1999). Rapid Avr9- and Cf-9 -dependent activation of MAP kinases in tobacco cell cultures and leaves: convergence of resistance gene, elicitor, wound, and salicylate responses. Plant Cell **11,** 273-287.

Romera, F.J., Alcantara, E. (1994). Iron-Deficiency Stress Responses in Cucumber (*Cucumis sativus* L.) Roots (A Possible Role for Ethylene?). Plant Physiol. **105,** 1133-1138.

Sablowski, R.W.M., Meyerowitz, E.M. (1998). A homolog of NO APICAL MERISTEM is an immediate target of the floral homeotic genes PETELA3/PISTILLATA. Cell **92,** 93-103.

Sakihama, Y., Cohen, M.F., Grace, S.C., Yamasaki, H. (2002). Plant phenolic antioxidant and prooxidant activities: phenolics-induced oxidative damage mediated by metals in plants. Toxicology **177,** 67-80.

Salt, D.E., Rauser, W.E. (1995). MgATP-Dependent Transport of Phytochelatins Across the Tonoplast of Oat Roots. Plant Physiol. **107,** 1293-1301.

Samaj, J., Ovecka, M., Hlavacka, A., Lecourieux, F., Meskiene, I., Lichtscheidl, I., Lenart, P., Salaj, J., Volkmann, D., Bogre, L., Baluska, F., Hirt, H. (2003). Involvement of MAP kinase SIMK and actin cytoskeleton in the regulation of root hair tip growth. Cell Biol. Int. **27,** 257-259.

Sambrook, J., Fritsch, E.F., Maniatis, T. (1989). Molecular Cloning: A Labotory Manual. (Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press).

Samuel, M.A., Ellis, B.E. (2002). Double jeopardy: both overexpression and suppression of a redox-activated plant mitogen-activated protein kinase render tobacco plants ozone sensitive. Plant Cell **14,** 2059-2069.

Sancenon, V., Puig, S., Mira, H., Thiele, D.J., Penarrubia, L. (2003). Identification of a copper transporter family in *Arabidopsis thaliana*. Plant Mol. Biol. **51,** 577-587.

Sanchez-Fernandez, R., Emyr Davies, T.G., Coleman, J.O.D., Rea, P.A. (2001). The *Arabidopsis thailana* ABC Protein Superfamily, a complete Inventory. J. Biol. Chem. **276,** 30231-30244.

Sanger, F., Nicklen, S., Coulsen, A.R. (1977). DNA Sequencing with chain-terminating inhibitors. Proc. Natl. Acad. Sci. USA **74,** 5463-5467.

Sangwan, V., Orvar, B.L., Beyerly, J., Hirt, H., Dhindsa, R.S. (2002). Opposite changes in membrane fluidity mimic cold and heat stress activation of distinct plant MAP kinase pathways. Plant J. **31,** 629-638.

Sanita di Toppi, L., Gabbrielli, R. (1999). Response to cadmium in higher plants. Environ. Exp. Bot. **41,** 105-130.

Sano, T., Nagata, T. (2002). The possible involvement of a phosphate-induced transcription factor encoded by phi-2 gene from tobacco in ABA-signaling pathways. Plant Cell Physiol. **43,** 12-20.

Sargent, T.D. (1987). Isolation of differentially expressed genes. Meth. Enzymol. **152,** 423-432.

Sarret, G., Saumitou-Laprade, P., Bert, V., Proux, O., Hazemann, J.L., Traverse, A.S., Marcus, M.A., Manceau, A. (2002). Forms of zinc accumulated in the hyperaccumulator *Arabidopsis halleri*. Plant Physiol. **130,** 1815-1826.

Schaaf, G., Catoni, E., Fitz, M., Schwacke, R., Schneider, A., von Wiren, N., Frommer, W.B. (2002). A putative role for the vacuolar calcium/manganese proton antiporter AtCAX2 in heavy metal detoxification. Plant Biol. **4,** 612-618.

Schaeffer, H.J., Weber, M.J. (1999). Mitogen-activated protein kinases: specific messages from ubiquitous messengers. Mol. Cell Biol. **19,** 2435-2444.

Schat, H., Liugany, M., Vooijs, R., Hartley-Whitaker, J., Bleeker, P.M. (2002). The role of phytochelatins in constitutive and adaptive heavy metal tolerances in hyperaccumulator and non-hyperaccumulator metallophytes. J. Exp. Bot. **53,** 2381-2392.

Schellenberger, A. (1989). Enzymkatalyse. (VEB Gustav Fischer Verlag Jena).

Schenk, H., Klein, M., Erdbrugger, W., Droge, W., Schulze-Osthoff, K. (1994). Distinct effects of thioredoxin and antioxidants on the activation of transcription factors NF-kappa B and AP-1. Proc. Natl. Acad. Sci. USA **91,** 1672-1676.

Schützendübel, A., Polle, A. (2002). Plant responses to abiotic stresses: heavy metal-induced oxidative stress and protection by mycorrhization. J. Exp. Bot. **53,** 1351-1365.

Seki, M., Narusaka, M., Ishida, J., Nanjo, T., Fujita, M., Oono, Y., Kamiya, A., Nakajima, M., Enju, A., Sakurai, T., Satou, M., Akiyama, K., Taji, T., Yamaguchi-Shinozaki, K., Carninci, P., Kawai, J., Hayashizaki, Y., Shinozaki, K. (2002). Monitoring the expression profiles of 7000 *Arabidopsis* genes under drought, cold and high-salinity stresses using a full-length cDNA microarray. Plant J. **31,** 279-292.

Seo, S., Sano, H., Ohashi, Y. (1999). Jasmonate-based wound signal transduction requires activation of WIPK, a tobacco mitogen-activated protein kinase. Plant Cell **11,** 289-298.

Shah, K., Dubey, R.S. (1995). Effect of Cadmium on RNA Level as Well as Activity and Molecular-Forms of Ribonuclease in Growing Rice Seedlings. Plant Physiol. Biochem **33,** 577-584.

Sharma, R., Rensing, C., Rosen, B.P., Mitra, B. (2000). The ATP hydrolytic activity of purified ZntA, a Pb (II) /Cd (II) /Zn(II) -translocating ATPase from *Escherichia coli*. J. Biol. Chem. **275,** 3873-3878.

Sharma, Y.K., Davis, R.K. (1995). Isolation of a novel *Arabidopsis* ozone-induced cDNA by differential display. Plant Mol. Biol. **29,** 91-98.

Siberil, Y., Doireau, P., Gantet, P. (2001). Plant bZIP G-box binding factors - Modular structure and activation mechanisms. Eur. J. Biochem. **268,** 5655-5666.

Siedlecka, A., Baszynski, T. (1993). Inhibition of Electron Flow around Photosystem-I in Chloroplasts of Cd-Treated Maize Plants Is Due to Cd-Induced Iron-Deficiency. Physiol. Plant. **87,** 199-202.

Simones-Araujo, J.L., Rodrigues, R.L., de A. Gerhartd, L.B., Mondego, J.M.C., Alves-Ferreira, M., Rumjanek, N.G., Margis-Pinheiro, M. (2002). Identification of differentially expressed genes by cDNA-AFLP technique during heat stress in cowpea nodules. FEBS Lett. **25875,** 1-7.

Solioz, M., Vulpe, C. (1996). CPx-type ATPases: A class of p-type ATPases that pump heavy metals. Trends Biochem.Sci. **21,** 237-241.

Stankovic, B., Vian, A., Henry-Vian, C., Davies, E. (2000). Molecular cloning and characterization of a tomato cDNA encoding a systemically wound-inducible bZIP DNA-binding protein. Planta **212,** 60-66.

Staxen, I.I., Pical, C., Montgomery, L.T., Gray, J.E., Hetherington, A.M., McAinsh, M.R. (1999). Abscisic acid induces oscillations in guard-cell cytosolic free calcium that involve phosphoinositide-specific phospholipase C. Proc. Natl. Acad. Sci. USA **96,** 1779-1784.

Steffen, W., Linck, R.W. (1989). Multiple immunoblot:a sensitive technique to stain proteins and detect multiple antigens on a single two-dimensional replica. Electrophoresis **10,** 714-718.

Stephan, U.W., Schmidke, I., Pich, A. (1994). Phloem Translocation of Fe, Cu, Mn, and Zn in Ricinus Seedlings in Relation to the Concentrations of Nicotianamine, an Endogenous Chelator of Divalent Metal-Ions, in Different Seedling Parts. Plant Soil **165,** 181-188.

Stohs, S.J., Bagchi, D. (1995). Oxidative Mechanisms in the Toxicity of Metal-Ions. Free Radic. Biol. Med. **18,** 321-336.

Strathmann, A., Kuhlmann, M., Heinekamp, T., Droge-Laser, W. (2001). BZI-1 specifically heterodimerises with the tobacco bZIP transcription factors BZI-2, BZI-3/TBZF and BZI-4, and is functionally involved in flower development. Plant J. **28,** 397-408.

Sun, Y., Hegamyer, G., Colburn, N.H. (1994). Molecular cloning of five messenger RNAs differentially expressed in preneoplastic or neoplastic JB6 mouse epidermal cells: one is homologous to human tissue inhibitor of metalloproteinases-3. Cancer Res. **5,** 1139-1144.

Sunkar, R., Kaplan, B., Bouche, N., Arazi, T., Dolev, D., Talke, I.N., Maathuis, F.J.M., Sanders, D., Bouchez, D., Fromm, H. (2000). Expression of a truncated tobacco NtCBP4 channel in transgenic plants and disruption of the homologous *Arabidopsis CNGC1* gene confer Pb^{2+} tolerance. Plant J. **24,** 533-542.

Sussman, M.R., Amasino, R.M., Young, C.J., Krysan, P.J., Austin-Phillpis, S. (2000). The *Arabidopsis* knockout facility at the University of Wisconsin-Madison. Plant Physiol. **124,** 1465-1467.

Suzuki, K., Shinshi, H. (1995). Transient Activation and Tyrosine Phosphorylation of a Protein-Kinase in Tobacco Cells Treated with a Fungal Elicitor. Plant Cell **7,** 639-647.

Suzuki, N., Koizumi, N., Sano, H. (2001). Screening of cadmium-responsive genes in *Arabidopsis thaliana*. Plant Cell Environ. **24,** 1177-1188.

Suzuki, N., Yamaguchi, Y., Koizumi, N., Sano, H. (2002). Functional characterization of a heavy metal binding protein CdI19 from *Arabidopsis*. Plant J. **32,** 165-173.

Tabata, K., Kashiwagi, S., Mori, H., Ueguchi, C., Mizuno, T. (1997). Cloning of a cDNA encoding a putative metal-transporting P-type ATPase from *Arabidopsis thaliana*. Biochim. Biophys. Acta **1326,** 1-6.

Tähtiharju, S., Palva, T. (2001). Antisense inhibition of protein phosphatase 2C accelerates cold acclimation in *Arabidopsis thaliana.* Plant J. **26,** 461-470.

Taiz, L., Zeiger, E. (1999). Physiologie der Pflanzen. (Spektrum Akademischer Verlag Heidelberg, Berlin).

Takagi, M., Satofuka, H., Amano, S., Mizuno, H., Eguchi, Y., Hirata, K., Miyamoto, K., Fukui, K., Imanaka, T. (2002). Cellular toxicity of cadmium ions and their detoxification by heavy metal-specific plant peptides, phytochelatins, expressed in Mammalian cells. J. Biochem. (Tokyo) **131,** 233-239.

Takahashi, K., Isobe, M., Knight, M.R., Trewavas, A.J., Muto, S. (1997). Hypoosmotic Shock Induces Increases in Cytosolic Ca^{2+} in Tobacco Suspension-Culture Cells. Plant Physiol. **113,** 587-594.

Tena, G., Renaudin, J.P. (1998). Cytosolic acidification but not auxin at physiological concentration is an activator of MAP kinases in tobacco cells. Plant J. **16,** 173-182.

The Arabidopsis Initiative. (2000). Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. Nature **408,** 796-815.

Thiele, D.J. (1988). ACE1 regulates expression of the *Saccharomyces cerevisiae* metallothionein gene. Mol. Cell Biol. **8,** 2745-2752.

Thimm, O., Essigmann, B., Kloska, S., Altmann, T., and Buckhout, T.J. (2001). Response of *Arabidopsis* to iron deficiency stress as revealed by microarray analysis. Plant Physiol. **127,** 1030-1043.

Thomine, S., Wang, R., Ward, J.M., Crawford, N.M., Schroeder, J.I. (2000). Cadmium and iron transport by members of a plant metal transporter family in *Arabidopsis* with homology to Nramp genes. Proc. Natl. Acad. Sci. USA **97,** 4991-4996.

Tommasini, R., Vogt, E., Fromenteau, M., Hortensteiner, S., Matile, P., Amrhein, N., Martinoia, E. (1998). An ABC-transporter of *Arabidopsis thaliana* has both glutathione-conjugate and chlorophyll catabolite transport activity. Plant J. **13,** 773-780.

Töpfer, R., Schell, J., Steinbiss, H.H. (1987). Versatile cloning vectors for transient gene expression and direct gene transfer in plant cells. Nucl. Acids Res. **16,** 8725.

Trezzini, G.F., Horrichs, A., Somssich, I.E. (1993). Isolation of putative defense-related genes from *Arabidopsis thaliana* and expression in fungal elicitor-treated cells. Plant Mol. Biol. **2,** 385-389.

Tron, A.E., Bertoncini, C.W., Chan, R.L., Gonzalez, D.H. (2002). Redox regulation of plant homeodomain transcription factors. J. Biol. Chem. **277,** 34800-34807.

Ulm, R., Ichimura, K., Mizoguchi, T., Peck, S.C., Zhu, T., Wang, X., Shinozaki, K., Paszkowski, J. (2002). Distinct regulation of salinity and genotoxic stress responses by *Arabidopsis* MAP kinase phosphatase 1. Embo J. **21,** 6483-6493.

Urao, T., Yamaguchi-Shinozaki, K., Shinozaki, K. (2000). Two-component systems in plant signal transduction. Trends Plant Sci. **5,** 67-74.

Usami, S., Banno, H., Ito, Y., Nishihama, R., Machida, Y. (1995). Cutting activates a 46-kilodalton protein kinase in plants. Proc. Natl. Acad. Sci. USA **92,** 8660-8664.

Van Assche, F., Clijsters, H. (1986a). Inhibition of Photosynthesis in *Phaseolus vulgaris* by Treatment with Toxic Concentration of Zinc - Effect on Ribulose-1,5- Bisphosphate Carboxylase Oxygenase. J. Plant Physiol. **125,** 355-360.

Van Assche, F., Clijsters, H. (1986b). Inhibition of Photosynthesis in *Phaseolus vulgaris* by Treatment with Toxic Concentrations of Zinc - Effects on Electron- Transport and Photophosphorylation. Physiol. Plant **66,** 717-721.

van der Biezen, E.A., Juwana, H., Parker, J.E., Jones, J.D.G. (2000). cDNA-AFLP Display for the Isolation of *Peronospora parasitica* Genes Expressed During Infection in *Arabidopsis thaliana*. Mol. Plant Microbe Interact. **13,** 895-898.

van der Zaal, B.J., Neuteboom, L.W., Pinas, J.E., Chardonnens, A.N., Schat, H., Verkleij, J.A.C., Hooykaas, P.J.J. (1999). Overexpression of a novel *Arabidopsis* gene related to putative zinc-transporter genes from animals can lead to enhanced zinc resistance and accumulation. Plant Physiol. **119,** 1047-1055.

van Hoof, N., Hassinen, V.H., Hakvoort, H.W.J., Ballintijn, K.F., Schat, H., Verkleij, J.A.C., Ernst, W.H.O., Karenlampi, S.O., Tervahauta, A.I. (2001). Enhanced copper tolerance in *Silene vulgaris* (Moench) Garcke populations from copper mines is associated with increased transcript levels of a 2b-type metallothionein gene. Plant Physiol. **126,** 1519-1526.

van Vliet, C., Anderson, C.R., Cobbett, C.S. (1995). Copper-sensitive mutant of *Arabidopsis thaliana*. Plant Physiol. **109,** 871-878.

Varotto, C., Maiwald, D., Pesaresi, P., Jahns, P., Salamini, F., Leister, D. (2002). The metal ion transporter IRT1 is necessary for iron homeostasis and efficient photosynthesis in *Arabidopsis thaliana*. Plant J. **31,** 589-599.

Vatamaniuk, O.K., Bucher, E.A., Ward, J.T., Rea, P.A. (2001). A new pathway for heavy metal detoxification in animals. Phytochelatin synthase is required for cadmium tolerance in *Caenorhabditis elegans*. J. Biol. Chem. **276,** 20817-20820.

Vatamaniuk, O.K., Mari, S., Lu, Y.P., Rea, P.A. (1999). AtPCS1, a phytochelatin synthase from *Arabidopsis*: Isolation and *in vitro* reconstitution. Proc. Natl. Acad. Sci. USA **96,** 7110-7115.

Velculescu, V.E., Zhang, L., Vogelstein, B., Kinzler, K.W. (1995). Serial analysis of gene expression. Science **5235,** 484-487.

Verkleij, J.A.C., Schat, H. (1990). Mechanisms of metal tolerance in higher plants. In Heavy Metal Tolerance in Plants: Evolutionary Aspects., A.J. Shaw, (CRC Press), 179-193.

Vert, G., Grotz, N., Dedaldechamp, F., Gaymard, F., Guerinot, M.L., Briata, J.F., Curie, C. (2002). IRT1, an *Arabidopsis* transporter essential for iron uptake from the soil and for plant growth. Plant Cell **14,** 1223-1233.

Vidmar, J.J., Tagmount, A., Cathala, N., Touraine, B., Davidian, J.-C.E. (2000). Cloning and characterization of a root specific high-affinity sulfate transporter from *Arabidopsis thaliana*. FEBS Lett. **475,** 65-69.

Villanueva, A., Lozano, J., Morales, A., Lin, X., Deng, X., Hengartner, M.O., Kolesnick, R.N. (2001). jkk-1 and mek-1 regulate body movement coordination and response to heavy metals through jnk-1 in *Caenorhabditis elegans*. Embo J. **20,** 5114-5128.

Vögeli-Lange, R., Wagner, G.J. (1990). Subcellular localization of cadmium and cadmium-binding peptides in tabacco leaves: implication of transport function for cadmium-binding peptides. Plant Physiol. **92,** 1086-1093.

von Wiren, N., Klair, S., Bansal, S., Briat, J.F., Khodr, H., Shioiri, T., Leigh, R.A., Hider, R.C. (1999). Nicotianamine chelates both Fe-III and Fe-II. Implications for metal transport in plants. Plant Physiol. **119,** 1107-1114.

Vos, P., Hogers, R., Bleeker, M., Reijans, M., van de Lee, T., Hornes, M., Frijters, A., Pot, J., Peleman, J., Kuiper, M. (1995). AFLP: a new technique for DNA fingerprinting. Nucl. Acids Res. **23,** 4407-4414.

Vranova, E., Langebartels, C., Van Montagu, M., Inze, D., Van Camp, W. (2000). Oxidative stress, heat shock and drought differentially affect expression of a tobacco protein phosphatase 2C. J. Exp. Bot. **51,** 1763-1764.

Wagner, U., Edwards, R., Dixon, D.P., F., M. (2002). Probing the Diversity ot the *Atabidopsis* glutathinone S-Transferase Gene Family. Plant Mol. Biol. **49,** 515-532.

Wang, M.L., Belmonte, S., Kim, U., Dolan, M., Morris, J.W., Goodman, H.M. (1999). A cluster of ABA-regulated genes on *Arabidopsis thaliana* BAC TO7MO7. Genome Res. **9,** 325-333.

Weber, M., Harada, E., Vess, C., v. Roepenack-Lahaye, E., Clemens, S. (in Druck). Comparative microarray analysis of *Arabidopsis thaliana* and *Arabidopsis halleri* roots identifies nicotianamine synthase, a ZIP transporter and other genes as potential metal hyperaccumulation factors. Plant J.

Weigel, D., Ahn, J.H., Blazquez, M.A., Borevitz, J.O., Christensen, S.K., Fankhauser, C., Ferrandiz, C., Kardailsky, I., Malancharuvil, E.J., Neff, M.M., Nguyen, J.T., Sato, S., Wang, Z.Y., Xia, Y., Dixon, R.A., Harrison, M.J., Lamb, C.J., Yanofsky, M.F., Chory, J. (2000). Activation tagging in *Arabidopsis*. Plant Physiol. **122,** 1003-1013.

Welsh, J., Chada, K., Dalal, S.S., R., C., McClelland, M. (1992). Arbitrarily primed PCR fingerprinting of RNA. Nucl. Acids Res. **20,** 4965-4970.

White, J.A., Todd, J., Newman, T., Focks, N., Girke, T., de Ilarduya, O.M., Jaworski, J.G., Ohlrogge, J.B., Benning, C. (2000). A new set of *Arabidopsis* expressed sequence tags from developing seeds. The metabolic pathway from carbohydrates to seed oil. Plant Physiol. **124,** 1582-1594.

White, P.J., Bowen, H.C., Demidchik, V., Nichols, C., Davies, J.M. (2002). Genes for calcium-permeable channels in the plasma membrane of plant root cells. Biochim. Biophys. Acta **1564,** 299-309.

Whiting, S.N., De Souza, M.P., Terry, N. (2001). Rhizosphere bacteria mobilize Zn for hyperaccumulation by *Thlaspi caerulescens*. Environ. Sci. Technol. **35,** 3144-3150.

Widmann, C., Gibson, S., Jarpe, M.B., Johnson, G.L. (1999). Mitogen-activated protein kinase: Conservation of a three- kinase module from yeast to human. Physiol. Rev. **79,** 143-180.

Willekens, H., Chamnongpol, S., Davey, M., Schraudner, M., Langebartels, C., Van Montagu, M., Inze, D., Van Camp, W. (1997). Catalase is a sink for H_2O_2 and is indispensable for stress defense in C3 plants. Embo J. **16,** 4806-4816.

Williams, L.E., Pittman, J.K., Hall, J.L. (2000). Emerging mechanism for heavy metal transport in plants. Biochim. Biophys. Acta **1465,** 104-126.

Wilson, C., Voronin, V., Touraev, A., Vicente, O., Heberle-Bors, E. (1997). A developmentally regulated MAP kinase activated by hydration in tobacco pollen. Plant Cell **9,** 2093-2100.

Wintz, H., Vulpe, C. (2002). Plant copper chaperones. Biochem. Soc. Trans. **30,** 732-735.

Winzeler, E.A., Schena, M., Davis, R.W. (1999). Fluorescence-Based Expression Monitoring Using Microarrays. Meth. Enzymol. **306,** 3-18.

Woeste, K.E., Kieber, J.J. (2000). A strong loss-of-function mutation in RAN1 results in constitutive activation of the ethylene response pathway as well as a rosette-lethal phenotype. Plant Cell **12,** 443-455.

Wollgiehn, R., Neumann, D. (1995). Stress-Response of Tomato Cell-Cultures to Toxic Metals and Heat-Shock - Differences and Similarities. J. Plant Physiol. **146,** 736-742.

Wortman, J.R., Haas, B.J., Hannick, L.I., Smith, R.K., Jr., Maiti, R., Ronning, C.M., Chan, A.P., Yu, C., Ayele, M., Whitelaw, C.A., White, O.R., Town, C.D. (2003). Annotation of the *Arabidopsis* genome. Plant Physiol. **132,** 461-468.

Wu, A., Wemmie, J.A., Edgington, N.P., Goebl, M., Guevara, J.L., Moye-Rowley, W.S. (1993). Yeast bZIP Proetins Mediate Pleiotropic Drug and Metal Resistance. J. Biol. Chem. **268,** 18850-18858.

Wu, Z., Liang, F., Hong, B., Young, J.C., Sussman, M.R., Harper, J.F., Sze, H. (2002). An endoplasmic reticulum-bound Ca^{2+}/Mn^{2+} pump, ECA1, supports plant growth and confers tolerance to Mn^{2+} stress. Plant Physiol. **130,** 128-137.

Xiang, C., Oliver, D.J. (1998). Glutathione metabolic genes coordinately respond to heavy metals and jasmonic acid in *Arabidopsis*. Plant Cell **10,** 1539-1550.

Xiang, C., Han, P., Lutziger, I., Wang, K., Oliver, D.J. (1999). A mini binary vector series for plant transformation. Plant Mol. Biol. **40,** 711-717.

Xiong, L., Schumaker, K.S., Zhu, J.K. (2002). Cell signaling during cold, drought, and salt stress. Plant Cell **14 Suppl,** S165-183.

Yablonski, D., Marbach, I., Levitzki, A. (1996). Dimerization of Ste5, a mitogen-activated protein kinase cascade scaffold protein, is required for signal transduction. Proc. Natl. Acad. Sci. USA **93,** 13864-13869.

Yamaguchi-Iwai, Y., Dancis, A., Klausner, R.D. (1995). AFT1: a mediator of iron regulated transcriptional control in *Saccharomyces cerevisiae*. Embo J. **14,** 1231-1239.

Yamaguchi-Iwai, Y., Ueta, R., Fukunaka, A., Sasaki, R. (2002). Subcellular localization of Aft1 transcription factor responds to iron status in *Saccharomyces cerevisiae*. J. Biol. Chem. **277,** 18914-18918.

Yamaguchi-Shinozaki, K., Shinozaki, K. (1994). A novel cis-acting element in an *Arabidopsis* gene is involved in responsiveness to drought, low-temperature, or high-salt stress. Plant Cell **6,** 251-264.

Yang, K.Y., Liu, Y.D., Zhang, S.Q. (2001). Activation of a mitogen-activated protein kinase pathway is involved in disease resistance in tobacco. Proc. Natl. Acad. Sci. USA **98,** 741-746.

Zenk, M.H. (1996). Heavy metal detoxification in higher plants - A review. Gene **179,** 21-30.

Zettl, R., Schell, J., Palme, K. (1994). Photoaffinity labeling of *Arabidopsis thaliana* plasma membrane vesicles by 5-azido-[7-3H]indole-3-acetic acid: identification of a glutathione S-transferase. Proc. Natl. Acad. Sci. USA **91,** 689-693.

Zhang, B., Foley, R.C., Singh, K.B. (1993). Isolation and Characterization of 2 Related *Arabidopsis* Ocs-Element bZIP Binding-Proteins. Plant J. **4,** 711-716.

Zhang, F., Romheld, V., Marschner, H. (1989). Effect of Zinc-Deficiency in Wheat on the Release of Zinc and Iron Mobilizing Root Exudates. Zeitung für Pflanzen- und Bodenkunde **152,** 205-210.

Zhang, S., Klessig, D.F. (1997). Salicylic acid activates a 48-kD MAP kinase in tobacco. Plant Cell **9,** 809-824.

Zhang, S., Klessig, D.F. (1998a). The tobacco wounding-activated mitogen-activated protein kinase is encoded by SIPK. Proc. Natl. Acad. Sci. USA **95,** 7225-7230.

Zhang, S., Klessig, D.F. (1998b). Resistance gene N-mediated de novo synthesis and activation of a tobacco mitogen-activated protein kinase by tobacco mosaic virus infection. Proc. Natl. Acad. Sci. USA **95,** 7433-7438.

Zhao, F.J., Lombi, E., Breedon, T., McGrath, S.P. (2000). Zinc hyperaccumulation and cellular distribution in *Arabidopsis halleri*. Plant Cell Environ. **23,** 507-514.

Zhao, H., Eide, D.J. (1997). Zap1p, a metalloregulatory protein involved in zinc-responsive transcriptional regulation in *Saccharomyces cerevisiae*. Mol. Cell Biol. **17,** 5044-5052.

Zhou, J.M., Goldsbrough, P.B. (1994). Functional Homologs of Fungal Metallothionein Genes from *Arabidopsis*. Plant Cell **6,** 875-884.

Zhu, H.N., Shipp, E., Sanchez, R.J., Liba, A., Stine, J.E., Hart, P.J., Gralla, E.B., Nersissian, A.M., Valentine, J.S. (2000). Cobalt binding to human and tomato copper chaperone for superoxide dismutase: Implications for the metal ion transfer mechanism. Biochemistry **39,** 5413-5421.

Zhu, J.K. (2001). Plant salt tolerance. Trends Plant Sci. **6,** 66-71.

Zhu, T., Budworth, P., B., H., Brown, D., Chang, H.-S., Zou, G., Wang, X. (2001). Toward elucidating the global gene expression patterns of developing *Arabidopsis*: Parallel analysis of 8300 genes by a high-density oligonucleotide probe array. Plant Physiol. Biochem. **39,** 221-242.